

Shape coexistence and configuration mixing in the neutron-deficient Hg isotopes

K. Nomura,^{1,*} R. Rodríguez-Guzmán,^{2,3} and L. M. Robledo⁴

¹*Institut für Kernphysik, Universität zu Köln, D-50937 Köln, Germany*

²*Department of Physics and Astronomy, Rice University, Houston, Texas 77005, USA*

³*Department of Chemistry, Rice University, Houston, Texas 77005, USA*

⁴*Departamento de Física Teórica, Universidad Autónoma de Madrid, E-28049 Madrid, Spain*

(Dated: December 3, 2012)

The coexistence of prolate and oblate geometrical shapes in the neutron-deficient mercury isotopes is analyzed on the basis of the Gogny energy density functional. Level energies and transition rates are calculated within the interacting boson model (IBM) with configuration mixing, whose parameters are determined by mapping the potential energy surface, obtained from the constrained self-consistent mean-field calculation employing the Gogny D1M functional, onto the IBM analog. The triaxial degree of freedom turns out to play an important role in the description of the shape coexistence and the configuration mixing in the considered mercury nuclei. An optimal choice of the configuration mixing IBM Hamiltonian is suggested, that meets the indication of the Gogny-D1M energy surface. The collective low-lying structure in the considered mercury chain is described nicely without adjustment, in particular the oblate-prolate band structures of the near mid-shell nuclei $^{184-188}\text{Hg}$.

PACS numbers: 21.10.Re, 21.60.Ev, 21.60.Fw, 21.60.Jz

I. INTRODUCTION

Both shape transition and shape coexistence in finite nuclei have been a theme of major interests in the field of low-energy nuclear structure study. In some nuclei, significantly low-lying excited 0^+ states close in energy to the 0^+ ground-state have been observed, that reveal the coexistence of different intrinsic shapes. Numerous efforts have already been made to better understand the nature of such stunning shape phenomena from both the theoretical and experimental sides (see Refs. [1–5] for review).

In terms of the nuclear shell model [6–11], the emergence of low-lying excited 0^+ states can be traced back to multiparticle-multihole excitations. This scenario applies to the neutron-deficient nuclei in the Lead region with $Z \approx 82$. In this case, two or four protons are excited from the $Z = 50-82$ major shell across the $Z = 82$ closed shell to the $h_{9/2}$ orbit. The residual interaction between the valence protons and neutrons becomes subsequently enhanced, leading to the lowering of the excited 0^+ energies. This effect is most significant around the neutron mid-shell $N = 104$.

In the neutron-deficient Hg isotopes, the weakly deformed oblate ground-state band coexists with a more deformed prolate one. Empirically, in these isotopes the oblate ground-state band stays constant in energy as a function of the neutron number down to ^{188}Hg , where it is crossed by intruder prolate configurations. The band-head of the intruder prolate band, i.e., the second 0^+ state, reaches a minimum in energy around ^{182}Hg and then goes up from ^{180}Hg [12]. From the experimental

point of view, this picture is quite vividly observed in the parabolic trend of the states belonging to the intruder prolate band as functions of the neutron number in the Hg chain. Given the recent advances in the experimental studies on the neutron-deficient Hg isotopes [12–15], it is quite timely, as well as significant, to pursue a theoretical description of the relevant spectroscopy that can be compared with available data.

Let us stress, that these heavy mass systems are currently beyond the reach of large-scale shell-model studies with a realistic nucleon-nucleon interaction. Therefore a drastic truncation scheme is required to make the problem more feasible. Such a framework is provided by the interacting boson model (IBM) [16], which employs the monopole s and quadrupole d bosons, associated to the $J = 0^+$ and 2^+ collective pairs of valence nucleons, respectively [17–19]. The collective levels and the transition rates are generated by the diagonalization of the boson Hamiltonian composed of only a few essential interaction terms. To handle the configuration mixing in the IBM framework, Duval and Barrett proposed to extend the boson Hilbert space to the direct sum of the configuration sub-spaces corresponding to the $2np-2nh$ excitation ($n = 0, 1, 2 \dots$) that comprises $N_B + 2n$ bosons [20]. Various features relevant to the shape coexistence phenomena in the Pb region have been investigated within the IBM configuration mixing model: the empirical collective structures from Po down to Pt isotopes [21–25], geometry and phases [26–28], and the algebraic features (in terms of the so called *intruder spin*) [29–31]. Nevertheless, one of the main difficulties was that, since too many parameters are involved in this prescription, one has to specify the form of the model Hamiltonian rather *a priori* and/or simply select the parameters by the fit to available data.

On the other hand, the energy density functional

*Electronic address: nomura@ikp.uni-koeln.de

(EDF) framework has been successful in the self-consistent mean-field study of bulk nuclear properties and collective excitations [32] with various classes of effective interactions, e.g., Skyrme [33, 34], Gogny [35], and those used within relativistic mean-field (RMF) models [36, 37]. The mean-field approximation provides the coexisting minima in the potential energy surface, which are associated to the different intrinsic geometrical shapes [38]. Deformation properties and collective excitations, relevant to the shape coexistence in the neutron-deficient Pb and Hg isotopes, have already been investigated using Skyrme [39, 40], Gogny [41–44] and the RMF interactions [45] as well as within the Nilsson-Strutinsky method [38].

Recently a method has been proposed that incorporates the merits of the EDF framework into the IBM by mapping the potential energy surface (PES) based on a given EDF onto the IBM system to generate, energy levels and transition rates with good symmetry quantum numbers [46]. The original idea of Ref. [46] has been extended to well deformed [47] and triaxial [48] systems and more recently, to the mixing of several multiparticle-multihole configurations in Lead isotopes [49] on the basis of the Duval-Barrett’s technique.

In this paper we apply the method of Ref. [49] to analyze the relevant spectroscopy related to the coexistence of different intrinsic shapes in the neutron-deficient $^{180-190}\text{Hg}$ isotopes. The predictive power of the model is also assessed by comparing its outcome with the available data. In addition, the optimal choice of the configuration mixing IBM Hamiltonian consistent with the EDF-based calculations is identified. Finally, the role of non-axial degrees of freedom in the microscopic EDF PES is stressed. We shall use the parametrization D1M of the Gogny-EDF [50], that has been shown (see, for instance [51–53]) to have a similar predictive power in the description of nuclear structure phenomena as the more conventional D1S [54] parameter set.

This paper is organized as follows: our theoretical framework is briefly summarized in Sec. II. Our results, for the microscopic (i.e., EDF) and the mapped PES in the considered Hg isotopes, the energy-level systematics, the detailed spectroscopy including E2 transitions, the spectroscopic quadrupole moment as well as the comparison with the phenomenological IBM are given in Secs. III and IV, respectively. Finally, Sec. V is devoted to the concluding remarks.

II. FRAMEWORK

We first perform a set of constrained Hartree-Fock-Bogoliubov (HFB) calculations using the Gogny-D1M [50] EDF to obtain the corresponding mean-field PES in terms of the geometrical quadrupole collective variables $q = (\beta, \gamma)$ [55]. Note that the PES denotes here the total mean-field energy, where neither mass parameter nor collective potential is considered explicitly, and that we only consider unprojected HFB PES.

Having the Gogny HFB PES, we subsequently map it onto the corresponding IBM PES. In order to treat the proton cross-shell excitation, we shall use the proton-neutron IBM (IBM-2) because it is more realistic than the original version of the IBM (IBM-1), which does not distinguish between proton and neutron degrees of freedom. The IBM-2 comprises neutron (proton) s_ν (s_π) and d_ν (d_π) bosons, which reflect the collective pairs of valence neutrons (protons) [19]. The number of neutron (proton) bosons, denoted as N_ν (N_π), equals half the number of the valence neutrons (protons). Since the Gogny-D1M PES exhibits two minima in the considered Hg nuclei, we take into account up to $2p-2h$ proton excitations. The doubly-magic nuclei ^{164}Pb and ^{208}Pb are regarded as boson vacua (inert cores) for the isotopes $^{180-190}\text{Hg}$. Therefore, N_π is fixed, $N_\pi = 1$ and 3 for the $0p-0h$ and the $2p-2h$ configurations, respectively. On the other hand, N_ν varies between 8 and 11.

The IBM Hamiltonian of the system comprising the normal ($0p-0h$) and the $2p-2h$ configurations is written as [20, 21]

$$\hat{H} = \hat{\mathcal{P}}_1 \hat{H}_1 \hat{\mathcal{P}}_1 + \hat{\mathcal{P}}_3 (\hat{H}_3 + \Delta) \hat{\mathcal{P}}_3 + \hat{H}_{\text{mix}}, \quad (1)$$

where $\hat{\mathcal{P}}_i$ ($i = 1, 3$) stands for the projection operator onto the $N_\pi = i$ configuration space. The operator $\hat{H}_i$ is the Hamiltonian for the configuration with $N_\pi = i$ bosons

$$\hat{H}_i = \epsilon_i \hat{n}_d + \kappa_i \hat{Q}_\nu^{\chi_\nu, i} \cdot \hat{Q}_\pi^{\chi_\pi, i} + \kappa'_i \hat{L} \cdot \hat{L} + \sum_{\rho' \neq \rho} \hat{V}_{\rho\rho'}, \quad (2)$$

where the first term $\hat{n}_d = \sum_\rho d_\rho^\dagger \cdot d_\rho$ ($\rho = \nu$ or π) stands for the d -boson number operator. The second term in Eq. (2) represents the quadrupole-quadrupole interaction between proton and neutron bosons with $\hat{Q}_{\rho, i}^{\chi_\rho, i} = d_\rho^\dagger s_\rho + s_\rho^\dagger \tilde{d}_\rho + \chi_{\rho, i} [d_\rho^\dagger \times \tilde{d}_\rho]^{(2)}$ being the quadrupole operator. The third term is relevant for rotationally deformed systems, with $\hat{L} = \sqrt{10} \sum_\rho [d_\rho^\dagger \times \tilde{d}_\rho]^{(1)}$ being the boson angular momentum operator. The fourth term on the right-hand side (RHS) of Eq. (2) stands for the three-body (cubic) boson term between proton and neutron bosons

$$\hat{V}_{\rho\rho\rho'} = \sum_L \kappa_{\rho\rho\rho', i}^{''(L)} [d_\rho^\dagger \times d_\rho^\dagger \times d_{\rho'}^\dagger]^{(L)} \cdot [\tilde{d}_{\rho'} \times \tilde{d}_\rho \times \tilde{d}_\rho]^{(L)}, \quad (3)$$

which is identified with the one used in Ref. [48]. For each $\rho = \nu$ and π , there are five linearly independent combinations in Eq. (3), identified by the value of $L = 0, 2, 3, 4$ and 6. In the present study, we only consider the $L = 3$ term, because its classical limit is proportional to $\cos^2 3\gamma$ the only term giving rise to a stable triaxial minimum at $\gamma \approx 30^\circ$ [48]. We have also considered the three-body interaction only between neutrons and protons since such proton-neutron correlation becomes more significant in medium-heavy and heavy nuclei. We assume $\kappa_{\pi\pi\nu, i}^{''(3)} = \kappa_{\pi\nu\nu, i}^{''(3)} \equiv \kappa_i''$, for simplicity [48].

In Eq. (1), Δ represents the energy off-set required to excite two protons from the $Z = 50 - 82$ to the $Z = 82 -$

126 major shells. In the same equation, the term $\hat{H}_{\text{mix}}$ stands for the interaction mixing between the normal and the $2p\text{-}2h$ configurations

$$\hat{H}_{\text{mix}} = \hat{\mathcal{P}}_3(\omega_s s_\pi^\dagger \cdot s_\pi^\dagger + \omega_d d_\pi^\dagger \cdot d_\pi^\dagger) \hat{\mathcal{P}}_1 + h.c., \quad (4)$$

where the parameters ω_s and ω_d are the mixing strength between the $N_\pi = 1$ and the $N_\pi = 3$ configurations.

It should be noted that, for each configuration, the Hamiltonian in Eq. (2) adopts the simplest possible form, with a minimal number of parameters consistent with the most relevant topology of the EDF PES. Up to the third term, the RHS of Eq. (2) is the standard form frequently used in a number of IBM-2 calculations [16]. In the present study, we include the so called $\hat{L} \cdot \hat{L}$ term since we have a relatively large number of bosons, which leads to a deformed rotational spectrum. On the other hand, as we will see below, the microscopic PES of the considered Hg nuclei exhibits a triaxial minimum, which requires the inclusion of the cubic term in the IBM Hamiltonian [48]. The physical significance of both the $\hat{L} \cdot \hat{L}$ and the cubic terms has been discussed in detail in Refs. [47] and [48], respectively.

The geometrical picture of a given IBM Hamiltonian is provided by the coherent-state framework [56]. Such a coherent state represents the boson intrinsic wave function specified by the deformation variables $\bar{q} = (\bar{\beta}_\nu, \bar{\beta}_\pi, \bar{\gamma}_\nu, \bar{\gamma}_\pi)$. One can take $\bar{\beta}_\nu = \bar{\beta}_\pi \equiv \bar{\beta}$ and $\bar{\gamma}_\nu = \bar{\gamma}_\pi \equiv \bar{\gamma}$ as only the isoscalar motion is of relevance in the following argument. The deformation $\bar{\beta}$ is assumed to be proportional to the one obtained within the HFB approximation while $\bar{\gamma}$ is taken the same in the HFB and IBM models [56].

In the configuration mixing model of IBM, one needs to consider the direct sum of the coherent state for the configuration with $N_{\pi,i} = i$ proton bosons [26], denoted here as $|\Phi_i(N_{\pi,i}, \beta, \gamma)\rangle$. The PES is obtained as the lower eigenvalue of the 2×2 coherent-state matrix [26]:

$$E(\beta, \gamma) = \begin{pmatrix} E_{11}(\beta, \gamma) & E_{31}(\beta) \\ E_{13}(\beta) & E_{33}(\beta, \gamma) + \Delta \end{pmatrix}. \quad (5)$$

The diagonal matrix element $E_{ii}(\beta, \gamma) = \langle \Phi_i(N_{\pi,i}, \beta, \gamma) | \hat{H}_i | \Phi_i(N_{\pi,i}, \beta, \gamma) \rangle$ on the RHS of Eq. (5) is given by

$$\begin{aligned} E_{ii}(\beta, \gamma) &= \frac{\epsilon'_i(N_\nu + N_{\pi,i})\bar{\beta}_i^2}{1 + \bar{\beta}_i^2} + \kappa_i N_\nu N_{\pi,i} \frac{\bar{\beta}_i^2}{(1 + \bar{\beta}_i^2)^2} \\ &\times \left[4 - 2\sqrt{\frac{2}{7}}(\chi_{\nu,i} + \chi_{\pi,i})\bar{\beta}_i \cos 3\gamma + \frac{2}{7}\chi_{\nu,i}\chi_{\pi,i}\bar{\beta}_i^2 \right] \\ &- \frac{1}{7}\kappa''_i N_\nu N_{\pi,i}(N_\nu + N_{\pi,i} - 2) \frac{\bar{\beta}_i^3}{(1 + \bar{\beta}_i^2)^3} \sin^2 3\gamma, \end{aligned} \quad (6)$$

where $\epsilon'_i = \epsilon_i + 6\kappa'_i$ and $\bar{\beta}_i \equiv C_{\beta,i}\beta$ with $C_{\beta,i}$ being the proportionality coefficient. The non-diagonal matrix element is given by $E_{ii'}(\beta, \gamma) =$

$\langle \Phi_{i'}(N_{\pi,i'}, \beta, \gamma) | \hat{H}_{\text{mix}} | \Phi_i(N_{\pi,i}, \beta, \gamma) \rangle$ ($i \neq i'$), and reads

$$\begin{aligned} E_{13}(\beta) &= E_{31}(\beta) \\ &= \sqrt{(N_{\pi,1} + 1)N_{\pi,3}} \left(\frac{\omega_s + \omega_d \bar{\beta}_3^2}{1 + \bar{\beta}_3^2} \right) \\ &\times \left(\frac{1 + \bar{\beta}_1 \bar{\beta}_3}{\sqrt{(1 + \bar{\beta}_1^2)(1 + \bar{\beta}_3^2)}} \right)^{N_\nu + N_{\pi,1}}. \end{aligned} \quad (7)$$

We also assume hereafter $\omega_s = \omega_d = \omega$ for simplicity.

The parameters ϵ'_i , κ_i , $\chi_{\nu,i}$, $\chi_{\pi,i}$, κ''_i , ω and Δ are determined through the same procedure as in Ref. [49], except that, in the present work, the Δ parameter determined from the PES is consistently used for the Hamiltonian [Eq. (1)] to be diagonalized. The $\hat{L} \cdot \hat{L}$ coefficient κ' is determined independently of the other parameters, that are derived from the PES. It is taken to match the Thouless-Valatin moment of inertia [57], calculated within the HFB approximation, to the one in the cranking approximation of the IBM-2 [58] (i.e., the one that is associated to each minimum on the PES [47, 59]).

Since the oblate HFB minimum occurs always at smaller deformation β (≈ 0.15) than the prolate one ($\beta \approx 0.3$) in the considered Hg isotopes, the Hamiltonians for the $0p\text{-}0h$ and $2p\text{-}2h$ configurations are associated to the oblate and the prolate minima, respectively. On the other hand, the locations of the oblate and the prolate minima on the β axis remain almost unchanged for all the considered nuclei. As a consequence, the scale factors $C_{\beta,i}$ remain almost constant for all nuclei, i.e., $C_{\beta,1} \approx 3$ and $C_{\beta,3} \approx 5$.

Finally, when the parameters are determined for an individual nucleus, the Hamiltonian in Eq. (1) is diagonalized in the enlarged model space in the boson m scheme. This gives rise to the energy spectra and wave functions for the excited states, that can be used to compute other spectral properties.

III. POTENTIAL ENERGY SURFACES

The Gogny-D1M and the mapped PES are plotted in Fig. 1 as functions of the $q = (\beta, \gamma)$ deformations. We have restricted the plots to configurations up to 3 MeV from the global minimum. Both microscopic and mapped PESs give β_2 values consistent with earlier calculations such as the Nilsson-Strutinsky method [38] and the collective model approach with the Gogny-D1S EDF [41], where $\beta_2 \approx -0.15$ and $0.25\text{-}0.3$ for the oblate and prolate configurations, respectively.

The Gogny-D1M PESs for $^{180,182}\text{Hg}$ display notable prolate minima at $\beta \approx 0.3$. On the other hand, the oblate minimum at $\beta \approx 0.15$ becomes gradually lower in energy when approaching ^{184}Hg , whose PES becomes softer along the γ direction. The prolate-oblate energy difference in the HFB PES reaches a minimum in ^{186}Hg which, at least within the Gogny-D1M HFB framework, represents the most dramatic example of shape coexistence in the considered Hg chain. Note that the lower




FIG. 1: (Color online) Microscopic (“D1M”) and mapped (“Mapped”) potential energy surfaces for the isotopes $^{180-190}\text{Hg}$ in the (β, γ) -plane are plotted up to 3 MeV from the absolute minimum. The microscopic results are obtained with the Gogny-D1M EDF.




FIG. 2: (Color online) Derived IBM parameters (a) ϵ_i , (b) κ_i , (c) $\chi_{\nu,i}$, (d) $\chi_{\pi,i}$, (e) κ''_i , (f) κ'_i , (g) Δ and (h) ω for the considered $^{180-190}\text{Hg}$ nuclei as functions of mass number A . Note that, in panel (f), the parameter κ'_i is plotted in keV unit. Figure legends in panels (a-f) are indicated in panel (a).

minimum becomes a bit off from the prolate axis with $\gamma \approx 5^\circ$. In ^{188}Hg , one can clearly see that the oblate

minimum becomes energetically favoured over the one around $\gamma \approx 10^\circ$ on the prolate side. In ^{190}Hg only the oblate minimum survives. The mapped IBM PES nicely follows the topology of the Gogny-D1M PES (i.e., locations of the minima as well as the curvatures around a given minimum in both the β and γ directions) for an individual nucleus. The overall systematic trend with neutron number is also well reproduced.

The derived IBM parameters [Eq. (1)] are depicted in Fig. 2 as functions of the mass number A . In agreement with its empirical boson number dependence [16], as well as with our previous findings [46], the single d -boson energy ϵ shown in panel a) exhibits a parabolic behavior with respect to the mid-shell. This also agrees, with the empirical evolution of the 2^+ excited state expected in a given isotope/isotone sequence. As can be observed, the ϵ_1 and ϵ_3 parameters, shown in the figure, roughly follow this empirical trend. Contrary to the fitting calculations [20, 21], the parameter ϵ_3 is always larger than ϵ_1 . The parameters $\kappa_{1,3}$, shown in panel b), do not change too much. Nevertheless, they are several times larger than the phenomenological ones ($\kappa_1 \approx -0.17 \sim -0.14$ MeV and $\kappa_3 \approx -0.14 \sim -0.11$ MeV) [22]. This is a direct consequence of the topology of the Gogny-D1M PES, whose energy minimum is too deep leading to a large curvature around the minimum on the β axis.

Generally, whether a given nucleus is prolate or oblate is specified by the sign of the sum $\chi_\nu + \chi_\pi$. As seen from Figs. 2(c) and 2(d), the sum has positive (negative) sign for the oblate (prolate) configurations, being consistent with the microscopic PES. In a previous study on Hg isotopes (e.g., [21]), the $2p\text{-}2h$ configuration has been considered in the rotational SU(3) limit of the IBM [16] by taking $\chi = \sqrt{7}/2 \approx 1.3$. The present result seems to contradict this assumption since both χ_ν and χ_π values for the $2p\text{-}2h$ configuration are away from the SU(3) limit of $\sqrt{7}/2$ reflecting the γ softness of the Gogny-D1M PES. Thus, not only the χ parameters but also the cubic term should be important for the description of the triaxiality in the considered Hg isotopes.

From Fig. 2(e), one sees that the derived κ_1'' value for both ^{186}Hg and ^{188}Hg is particularly large in agreement with the Gogny-D1M PES [Fig.1] of the two nuclei displaying the most notable γ softness in the considered isotopic chain. The $\hat{L} \cdot \hat{L}$ coefficient κ' , shown in panel f), appears to be almost constant, but its sign is of particular relevance here. The positive (negative) sign for the normal ($2p\text{-}2h$) configuration implies that the inclusion of the $\hat{L} \cdot \hat{L}$ term reduces (enlarges) the moment of inertia of the unperturbed collective band of the configuration. As we will show later (Sec. IV), the $\hat{L} \cdot \hat{L}$ may have a crucial impact on the level spacing for each band.

The energy offset Δ in Fig. 2(g) changes with neutron number symmetrically with respect to ^{184}Hg . The energy needed to excite two protons across the $Z = 82$ closed shell becomes maximal for this mid-shell nucleus because the intruder $2p\text{-}2h$ configuration gains maximal energy through deformation. Finally, as can be observed from




FIG. 3: (Color online) Level-energy systematics for $^{180-190}\text{Hg}$ isotopes with mass number. Theoretical level energies resulting from the mapped IBM-2 Hamiltonian based on the Gogny-D1M EDF (a) are compared with the experimental (b) energies taken from the ENSDF database [60] and from Ref. [12]. To guide the eye, dashed and solid lines connect the yrast and the non-yrast states, respectively.

panel h), the mixing strength ω remains rather constant

IV. SPECTRAL PROPERTIES

Let us now discuss how the topology, as well as the evolution, of the microscopic PES and the derived parameters correspond to the spectroscopic properties of the considered Hg isotopes.

A. Level energy systematics

We show in Fig. 3 the low-lying excited states of the isotopes $^{180-190}\text{Hg}$ with spins $J \leq 6^+$. The theoretical energy levels in Fig. 3(a), obtained through the diagonalization of the mapped IBM-2 Hamiltonian, are compared with the corresponding experimental data, shown in panel b), taken from the Evaluated Nuclear Structure Data File (ENSDF) [60] and from Ref. [12] for ^{180}Hg .

We observe, a reasonable qualitative agreement between the calculated and the experimental levels. The lowest two 0^+ states in the neutron-deficient Hg isotopes originate either from the oblate or the prolate configurations. Starting from ^{190}Hg down to the mid-shell nucleus ^{184}Hg , the calculated yrast states come down following the trend of the experimental levels. A rather good agreement with the experimental data has been obtained for Hg nuclei with $A \geq 184$, including the ordering of the energy levels and the location of the 0_2^+ state. Experimentally, the 0_2^+ excited state in $^{180,182}\text{Hg}$ (the bandhead of the intruder prolate band) is still lying close to, and almost degenerate with, the 2_1^+ excited state around the




FIG. 4: (Color online) Fraction of the intruder $2p\text{-}2h$ configuration in the wave functions of the $0_{1,2}^+$, the $2_{1,2}^+$ and the $4_{1,2}^+$ excited states of the considered $^{180-190}\text{Hg}$ isotopes (in %). The bars for each nucleus (from left to right) correspond to 0_1^+ , 0_2^+ , 2_1^+ , 2_2^+ , 4_1^+ and 4_2^+ excited states, respectively, as indicated in the figure.

middle of the major shell. On the other hand, the theoretical non-yrast levels increase their excitation energies for $A \leq 182$. In particular, the 0_2^+ excitation energy is too large in comparison with the corresponding experimental value. This can be due to the fact that the mean-field oblate-prolate energy difference predicted with the Gogny-D1M EDF could be too large for these two nuclei. In general, the HFB oblate-prolate energy difference should roughly correspond to the excitation energy of the 0_2^+ before mixing. Once the two configurations are mixed, the 0^+ excitation energy becomes more or less larger than the energy difference between the two mean-field minima. This is due to the level repulsion effect that is seen in $^{186,188}\text{Hg}$. However, the 0_2^+ excitation energy for $^{180,182,184}\text{Hg}$ seems to be almost equal to, and even smaller than, the oblate-prolate energy difference in the corresponding PES. The reason is probably that, mainly due to the triaxiality and to the shift of the d boson energy by $6\kappa'$ necessary to keep the topology of the PES, the unperturbed 0_2^+ excitation energy is sufficiently smaller than the oblate-prolate energy difference in the HFB PES. As a result, the energy of the 0_2^+ level after the configuration mixing, may be even smaller than the energy difference between the two mean-field minima.

In particular, to assess the effect of the cubic term on the spectrum, we performed a test calculation, in which the cubic term was not included in both configurations. Without the cubic term, the derived mixing strength ω turned out to be several tens per cent larger than the one in Fig. 2(h) in order to reproduce the topology of the EDF PES along the γ direction. This resulted in stronger level repulsion between the two 0^+ eigenstates, and thus the 0_2^+ excitation energy was overestimated by several hundred keV in comparison with Fig. 3(a).




FIG. 5: (Color online) Level scheme for the ^{188}Hg nucleus. The level energies and the $B(E2)$ values (numbers written along arrows) obtained from the mapped IBM-2 Hamiltonian based on the Gogny-D1M EDF are compared with the corresponding experimental data [60]. The oblate and the prolate bands are indicated by “Obl.” and “Prol.”, respectively. To help identify the corresponding values in the plot, $B(E2; 4_2^+ \rightarrow 2_2^+) = 84$ W.u. and $B(E2; 0_2^+ \rightarrow 2_2^+) = 850$ (in Weisskopf units) in the right panel.

B. Detailed level scheme

To identify band structures, we now analyze the detailed level scheme of individual nuclei, including both the inter- and the intra-band E2 transition rates. We will consider the isotopes $^{182,184,186,188}\text{Hg}$ near the neutron mid-shell $N \approx 104$. These isotopes exhibit, the most spectacular competition between different shapes among the considered Hg nuclei. To facilitate the comparison with the experimental data, the excited states shown below are classified into either oblate or prolate band according to the prolate-oblate predominance of the wave function of the state and/or the E2 transition sequence.

To this end, we have first computed the probability of the basis states in the wave function of the state of interest. In Fig. 4 we have plotted, the fraction of the $2p\text{-}2h$ component in the wave functions of the $0_{1,2}^+$, $2_{1,2}^+$ and the $4_{1,2}^+$ states. One sees that the yrast states in $^{180,182}\text{Hg}$ are composed of the intruder prolate configuration. The two configurations are mixed for each of the low-lying states of $^{184,186}\text{Hg}$. On the other hand, for the nucleus ^{188}Hg the tendency is the opposite to the one in ^{182}Hg . Only a small fraction of the intruder component plays a role in the low-lying states of ^{190}Hg . It appears that the excited 0^+ state of the nucleus ^{190}Hg is originated from a single oblate configuration, consistently with the empirical observation [41].

The $B(E2; J \rightarrow J')$ transition rate reads

$$B(E2; J \rightarrow J') = \frac{1}{2J+1} |\langle J' || \hat{T}^{(E2)} || J \rangle|^2, \quad (8)$$

where $|J\rangle$ ($|J'\rangle$) represents the wave function of the initial (final) state with spin J (J'). The E2 operator $\hat{T}^{(E2)}$




FIG. 6: (Color online) Same as Fig. 5, but for ^{186}Hg nucleus and, $E(3_2^+) = 1.970$, $E(4_1^+) = 1.959$ (in MeV), $B(E2; 2_2^+ \rightarrow 0_2^+) = 88$, $B(E2; 3_1^+ \rightarrow 2_3^+) = 136$, $B(E2; 3_2^+ \rightarrow 2_4^+) = 188$, $B(E2; 5_2^+ \rightarrow 3_2^+) = 30$, $B(E2; 5_2^+ \rightarrow 4_4^+) = 62$ and $B(E2; 7_1^+ \rightarrow 6_3^+) = 23$ (in W.u.) in the right panel.

is given as $\hat{T}^{(E2)} = \sum_{\rho,i=1,3} e_{\rho,i} \hat{P}_i \hat{Q}_{\rho}^{\chi_{\rho,i}} \hat{P}_i$, where $\hat{Q}_{\rho}^{\chi_{\rho,i}}$ is identified with the quadrupole operator in Eq. (2). We considered the same $\chi_{\rho,i}$ value as the one used in the diagonalization of the Hamiltonian in Eq. (1). This is equivalent to the so-called consistent-Q formalism in the IBM-1 [61]. The boson effective charge $e_{\rho,i}$ is assumed to be the same for protons and neutrons, i.e., $e_{\nu,i} = e_{\pi,i} \equiv e_i$. The e_1 value is fixed to $e_1 = 0.166$ eb, taken from Ref. [22], while the e_3 value is determined for each individual nucleus according to the empirical relation $e_3/e_1 \approx |\kappa_3/\kappa_1|$ [20, 22].

We show in Fig. 5 the detailed level scheme of the nucleus ^{188}Hg . The spectra and the $B(E2)$ transition strengths, computed from the mapped IBM-2 Hamiltonian based on the Gogny-D1M EDF, are compared with the experimental data in the same figure. A nice agreement between theory and experiment is obtained regarding the band structure, the energy of the 0_2^+ state as well as the level ordering. A pronounced mixing between the two configurations can be confirmed from the $B(E2)$ values for the transitions $6_{1,2}^+ \rightarrow 4_{1,2}^+$. They reflect the sizable amount of the mixing seen in the wave functions of the two 6^+ excited states [the wave function of the 6_1^+ (6_2^+) state consists of 69.1 (38.2)% of the $2p-2h$ component]. Mainly due to the uncertainty in the boson effective charges, the $B(E2)$ values for the transitions $2_2^+ \rightarrow 2_1^+$ and $2_2^+ \rightarrow 0_1^+$ are rather larger than the corresponding available data. On the other hand, the branching ratio for these transitions compare well with the experiment.

Consistent with the experimental evidence, the 0^+ ground-state in ^{186}Hg (see, Fig. 6) is predicted to be of oblate nature. The normal and the intruder configurations are substantially mixed in the wave functions of the

lowest two 0^+ states. According to Fig. 4, the 0_1^+ (0_2^+) state comprises 36.7 (62.3)% of the intruder $2p-2h$ configuration. Actually, the prolate configuration is quite notable in the mean-field PES shown in Fig. 1. In addition, our calculations seem to reproduce the experimental level scheme including the $B(E2)$ pattern: the oblate ground-state band is rather stretched, while the intruder prolate band built on the 0_2^+ excited state is compressed.

Such systematics in the level spacing for the lower band in each configuration, which is likely to be the $K^\pi = 0^+$ band, can only be reproduced with the inclusion of the $\hat{L} \cdot \hat{L}$ term. Actually, we have also performed a test calculation for all nuclei considered, that did not include the $\hat{L} \cdot \hat{L}$ term, and especially the intruder prolate band turned out to be, as expected, too stretched inconsistently with the data. This finding represents, of course, only a practical side of the $\hat{L} \cdot \hat{L}$ term. More importantly, this result implies that to describe the shape coexistence and hence the configuration mixing in the considered Hg isotopes within the present model, not only the static HFB PES but a specific dynamics should be also taken into account through the $\hat{L} \cdot \hat{L}$ term with nonzero angular frequency. In addition, a pronounced mixing between the two configurations is observed from the branching ratios of the E2 transition, such as for the $4_{1,2}^+ \rightarrow 2_{1,2}^+$ transitions, which are indeed comparable with each other.

We also address the study of collective bands, other than those built on the two 0^+ states, for ^{186}Hg as there are sufficient experimental data to compare with. From Fig. 6 one sees that the present calculation also reproduces the excitation energies with odd spin quite nicely. The odd-spin states, 3_1^+ , 5_1^+ and 7_1^+ (3_2^+ , 5_2^+ and 7_2^+), are predicted to be the members of the prolate (oblate) band. Due to the strong E2 transition sequence, it is




FIG. 7: (Color online) Same as Fig. 5, but for ^{184}Hg nucleus and $B(E2; 3_1^+ \rightarrow 2_3^+) = 153 \text{ W.u.}$

also likely that a set of the states 2_3^+ , 3_1^+ , 4_3^+ , 5_1^+ , 6_3^+ and 7_1^+ (2_4^+ , 3_2^+ , 4_4^+ , 5_2^+ , 6_4^+ and 7_2^+), forms a quasi- γ , i.e., $K^\pi = 2^+$, band for the prolate (oblate) configuration. These two quasi- γ bands seem to be close in energy, with 2^+ bandheads being within 400 keV. One can also observe in both quasi- γ bands quite strong E2 transitions from the 3^+ level to the corresponding 2^+ bandhead, which is in the same order of magnitude as the $2^+ \rightarrow 0^+$ E2 transition in each $K^\pi = 0^+$ band, and those between the members of the quasi- γ band. Note that this prediction (i.e., the existence of the two quasi- γ bands) is consistent with the empirical assignment of these levels [41], including the collective model description based on Gogny-D1S EDF. In addition, one should notice that the quasi- γ band in the prolate configuration looks similar to the one predicted within the rigid-triaxial rotor model of Davydov and Filippov, that is, the doubles $(2_\gamma^+, 3_\gamma^+)$, $(4_\gamma^+, 5_\gamma^+)$, $(6_\gamma^+, 7_\gamma^+)$, ... etc [62]. From the comparison with the IBM phenomenology for ^{186}Hg [22], one sees that the present result exhibits a similar level of agreement with the experimental one regarding the energy spectra of the oblate ground-state band with $J \leq 6^+$. Our result also better reproduces the 3_1^+ state while, as addressed in Sec. IV A, the prolate 0_2^+ band-head energy is a bit more overestimated.

From Fig. 7, one observes that the quality of agreement with the experiment in ^{184}Hg is similar to the one obtained for ^{186}Hg . From the strong E2 transitions to the corresponding 0^+ bandheads, we assume that the 2_1^+ and the 2_2^+ states belong to the oblate and the prolate bands, respectively. As seen from Fig. 4, although the wave function of the 2_1^+ (2_2^+) state is predominantly of the $2p-2h$ ($0p-0h$) nature, the mixing between the two configurations is quite strong: 65.4 (37.5) % of the 2_1^+ (2_2^+) wave function is comprised of the intruder $2p-2h$ configuration. Again, the strong mixing between the two




FIG. 8: (Color online) Same as Fig. 5, but for ^{182}Hg nucleus.

configurations is confirmed by the branching ratios of the $4_{1,2}^+$ to the $2_{1,2}^+$ excited states. Similar to ^{186}Hg , we observe a significant role played by the $\hat{L} \cdot \hat{L}$ term in the description of the level spacing. As seen in Fig. 7, the present study predicts the existence of the two quasi- γ bands for ^{184}Hg as well. It should be of interest to identify such band structure experimentally. In the present framework, such systematics of the γ band seems to be seen most vividly in $^{184,186}\text{Hg}$ since the oblate and/or the prolate mean-field minimum is the softest and the closest in energy among the considered Hg isotopes. Regarding the nucleus ^{188}Hg , however, any sequence of the quasi- γ band structure has not been obtained in our calculations (not shown in Fig. 5). This agrees well with available data [60]. On the other hand, the predicted 3_1^+ energy (1.437 MeV) agrees well with the data (1.455 MeV) [60].

In Fig. 8, we show the results corresponding to the nucleus ^{182}Hg . Rather contrary to the experimental assign-




FIG. 9: (Color online) Calculated spectroscopic quadrupole moments $Q^{(s)}$ for the lowest two excited 2^+ states of the considered Hg nuclei as functions of mass number.

ment, the present calculation predicts the ground-state band to have an intruder prolate nature, as 75.4% of the 0_1^+ state is dominated by the $2p-2h$ configuration (see, Fig. 4). A clear collective pattern is seen from the calculated E2 transition sequence for both the prolate and the oblate bands, although the $B(E2; 2_1^+ \rightarrow 0_1^+)$ value is rather overestimated. Another major deviation from the experimental data occurs in the intra-band transition from the 4_2^+ state in the oblate band to the 2_1^+ state in the prolate band. The corresponding experimental value, $B(E2; 4_1^+ \rightarrow 2_1^+) = 176(19)$ W.u., is quite large in comparison to the $2_1^+ \rightarrow 0_1^+$ E2 transition strength reflecting the very strong mixing between the different configurations. In our calculations the intra-band transition $B(E2; 4_2^+ \rightarrow 2_1^+) = 2$ W.u., while $B(E2; 2_1^+ \rightarrow 0_1^+) = 116$ W.u., although there exists a certain amount of mixing between the two configurations in the corresponding eigenstates (see, Fig. 4). A possible reason for this deviation could be that, in comparison to the experimental data, the theoretical 0_2^+ excitation energy is too large, leading to a weak E2 transition. This problem arises mainly because the energy difference between the oblate and the prolate minima is too large in the HFB PES (see, Fig. 1). A similar argument holds for ^{180}Hg , whose wave functions for the low-lying states look very similar in the composition of the two configurations to the ones for ^{182}Hg in Fig. 4, exception made of the 4_1^+ state.

C. Spectroscopic quadrupole moment

In order to better understand the deformation properties of the considered Hg isotopes, we have also analyzed the spectroscopic quadrupole moments for the lowest two 2^+ excited states belonging either to the first oblate or prolate band. The spectroscopic quadrupole

moment $Q^{(s)}$ for a state with spin J reads

$$Q^{(s)}(J) = \sqrt{\frac{16\pi}{5}} \begin{pmatrix} J & 2 & J \\ -J & 0 & J \end{pmatrix} \langle J || \hat{T}^{(E2)} || J \rangle. \quad (9)$$

The calculated quadrupole moments for the 2_1^+ and the 2_2^+ states in the $^{180-190}\text{Hg}$ nuclei, are shown in Fig. 9. For the lighter isotopes $^{180-184}\text{Hg}$, $Q^{(s)}(2_1^+)$ [$Q^{(s)}(2_2^+)$] is negative (positive), reflecting the prolate (oblate) nature of the state. On the other hand, $Q^{(s)}(2_1^+)$ becomes positive for $A \geq 186$. For both $^{184,186}\text{Hg}$, the predicted $Q^{(s)}$ values are smaller in magnitude than in other nuclei, as there is significant triaxiality in these nuclei. The overall systematic trend in $Q^{(s)}(2_{1,2}^+)$ seems to correlate well with the evolution of mean-field minima shown in Fig. 1 and with the structure of the corresponding wave functions in Fig. 4.

To compare directly with the experimental data, one can also deduce the intrinsic deformation parameter $\beta_2(J)$ from $Q^{(s)}(J)$, through the relation

$$\beta_2(J) = \sqrt{\frac{5}{16\pi}} \frac{4\pi Q^{(i)}(J)}{3ZR^2}, \quad (10)$$

where $R = 1.2A^{1/3}$ fm and $Q^{(i)}(J)$ represents the intrinsic quadrupole moment given by

$$Q^{(i)}(J) = -\frac{2J+3}{J} Q^{(s)}(J) \quad (11)$$

with the assumption $K = 0$. In the present work, the calculated $\beta_2^{obl}(2^+)$ value is generally consistent with the one taken from the mean-field PES in Fig. 1 ($\beta_2 \approx 0.15$), whereas the $\beta_2^{prol}(2^+)$ one is considerably smaller. In fact, for $^{180,182,188,190}\text{Hg}$, we obtain $-0.17 \leq \beta_2^{obl}(2^+) \leq -0.15$ and $0.15 \leq \beta_2^{prol}(2^+) \leq 0.16$, while for $^{184,186}\text{Hg}$ both β_2^{prol} and $|\beta_2^{obl}|$ are much smaller than 0.1. These small $\beta_2(2^+)$ values seem to be rather reasonable because the effects of the configuration mixing and the non-axial degrees of freedom (characterized by the rather small magnitudes of χ_ν and χ_π), which we have emphasized so far, are taken into account through the diagonalization.

D. Comparison with the phenomenological fit

Finally, we assess the validity of the set of IBM parameters derived here by comparing it to the one in Ref. [21] determined from a fit to the experimental spectra. Both sets of parameters lead to reasonable agreement with the data, but they differ in their description of the ground-state shape.

To illustrate this, we show in Fig. 10 the PES for the nucleus ^{184}Hg using the set of parameters used in Ref.[21]. The IBM Hamiltonian for both configurations in Ref.[21] was identical to the one on the RHS of Eq. (2) up to the second term, but included the neutron-boson




FIG. 10: (Color online) The potential energy surface of the ^{184}Hg nucleus, based on the parameters determined from a phenomenological fit [21] to the experimental spectra. The PES is drawn in terms of the boson quadrupole deformation variables $\bar{q} = (\bar{\beta}, \bar{\gamma})$.

interaction term $\hat{V}_{\nu\nu} = \frac{1}{2} \sum_{L=0,2,4} C_L^\nu (d_\nu^\dagger \times d_\nu^\dagger)^{(L)} \cdot (\tilde{d}_\nu \times \tilde{d}_\nu)^{(L)}$. Note that the PES in Fig. 10 is plotted using the same energy scale as in Fig. 1 but in terms of the boson variables $\bar{q} = (\bar{\beta}, \bar{\gamma})$, as they are not connected to the geometrical deformation variables.

One sees in Fig. 10 the spherical minimum, which leads to the vibrational U(5) [16] spectra, as well as the second minimum on the prolate axis, corresponding to the intruder deformed band equivalent to the SU(3) limit of the IBM [16]. Therefore, the corresponding energy spectrum can be considered as the coupling of the two limits. However, this picture seems to be rather simplistic. If one compares the mapped PES in Fig. 1 and the one in Fig. 10 a difference is observed in the role of triaxiality and the pronounced oblate minimum at $\beta \approx 0.15$ of the mapped PES. Indeed, the mapped PES is generally rather sharp in both the oblate and prolate axes and exhibits a richer structure along the γ direction. On the other hand, the phenomenological PES appears to be flat in both β and γ directions.

A possible reason for the difference would be the limited number and/or species of bosons in the IBM. Regarding the former, an inert core is assumed in the boson intrinsic (coherent) state and so the boson number is already a good quantum number. On the other hand, within the self-consistent EDF calculation carried out here the model space consists of all nucleons but the particle number is still not a good quantum number. In the mapped PES, such effect may be renormalized into the parameters in the configuration-mixing IBM Hamiltonian through the calibration with the EDF PES and so does not matter too much in the present framework. In this respect, it should be also interesting to see how qualitatively/quantitatively could a symmetry-projected calculation change the topology of the mean-field PES and the resulting spectrum in the considered Hg nuclei.

V. SUMMARY

The shape coexistence phenomena in the neutron-deficient Hg isotopes has been investigated within the

IBM-2 with configuration mixing based on the Gogny-D1M EDF. The parameters of the IBM Hamiltonian were determined through the mapping of the Gogny-D1M PES onto the one provided by an IBM-2 Hamiltonian in the boson condensate.

The importance of the γ degree of freedom has been emphasized in the description of the ground-state shape of the considered Hg nuclei with the Gogny-D1M EDF. Spectroscopic quantities that help to identify the oblate-prolate shape coexistence, resulting from the mapped Hamiltonian, have been compared with the experimental data. The mapped Hamiltonian provides us with a fairly good description of the observed energy levels without any adjustment, including the predominance of the oblate or the prolate configuration in each collective band. Particularly in the near mid-shell nuclei $^{184,186,188}\text{Hg}$, both the calculated oblate and prolate bands indicate a notable collective pattern in the in-band and the intra-band E2 transition rates, the latter being a strong evidence for the mixing between the two configurations.

It has been shown that the observed collective band structure for both the oblate and the prolate configurations is reproduced only with the proposed Hamiltonian including the cubic term and the $\hat{L} \cdot \hat{L}$ term. The cubic term has been introduced following the indications of the EDF PES, which reveals the importance of the triaxial degrees of freedom. It is also necessary not only to use the static HFB PES information, but also to consider the response of the system to nonzero angular frequency as required by the $\hat{L} \cdot \hat{L}$ term. In this respect, the present work suggests a possible optimal choice of the configuration mixing IBM Hamiltonian as well as the necessity to take into account the non-axial degrees of freedom within an EDF framework for the neutron-deficient Hg isotopes.

Nevertheless, the disagreement between our results and the data in $^{180,182}\text{Hg}$ should not be overlooked. This problem has been mainly attributed to the parametrization and/or the form of the considered EDF, which predicts that the two HFB minima differ too much in energy. An extension of the present EDF framework would help to clarify this problem.

Acknowledgments

Authors would like to thank P. von Brentano, M. Hackstein, J. Jolie, T. Otsuka and N. Shimizu for useful discussions. Author K. N. acknowledges the support of the JSPS Postdoctoral Fellowships for Research Abroad. L. M. R. acknowledges support of MINECO through grants Nos. FPA2009-08958 and FIS2009-07277 as well as the Consolider-Ingenio 2010 program CPAN CSD2007-00042 and MULTIDARK CSD2009-00064.

-
- [1] K. Heyde, P. Van Isacker, M. Waroquier, J. L. Wood, and R. A. Meyer, *Phys. Rep.* **102**, 291 (1983).
- [2] J. L. Wood, K. Heyde, W. Nazarewicz, M. Huyse, and P. van Duppen, *Phys. Rep.* **215**, 101 (1992).
- [3] A. N. Andreyev, M. Huyse, P. Van Duppen, L. Weissman, D. Ackermann, J. Gerl, F. P. Hessberger, S. Hofmann, A. Kleinbühl, G. Münzenberg, et al., *Nature (London)* **405**, 430 (2000).
- [4] R. Julin, K. Helariutta, and M. Muikku, *J. Phys. G* **27**, R109 (2001).
- [5] K. Heyde and J. L. Wood, *Rev. Mod. Phys.* **83**, 1467 (2011).
- [6] P. Federman and S. Pittel, *Phys. Lett. B* **69**, 385 (1977).
- [7] P. Van Duppen, E. Coenen, K. Deneffe, M. Huyse, K. Heyde, and P. Van Isacker, *Phys. Rev. Lett.* **52**, 1974 (1984).
- [8] K. Heyde, P. Van Isacker, R. F. Casten, and J. L. Wood, *Phys. Lett. B* **155**, 303 (1985).
- [9] K. Heyde, J. Jolie, J. Moreau, J. Ryckebusch, M. Waroquier, P. Van Duppen, M. Huyse, and J. L. Wood, *Nucl. Phys. A* **466**, 189 (1987).
- [10] K. Heyde and R. A. Meyer, *Phys. Rev. C* **37**, 2170 (1988).
- [11] K. Heyde, J. Jolie, H. Lehmann, C. De Coster, and J. L. Wood, *Nucl. Phys. A* **586**, 1 (1995).
- [12] J. Elseviers, A. N. Andreyev, S. Antalic, A. Barzakh, N. Bree, T. E. Cocolios, V. F. Comas, J. Diriken, D. Fedorov, V. N. Fedosseyev, et al., *Phys. Rev. C* **84**, 034307 (2011).
- [13] T. Grahn, A. Petts, M. Scheck, P. A. Butler, A. Dewald, M. B. G. Hornillos, P. T. Greenlees, A. Görden, K. Helariutta, J. Jolie, et al., *Phys. Rev. C* **80**, 014324 (2009).
- [14] M. Scheck, T. Grahn, A. Petts, P. A. Butler, A. Dewald, L. P. Gaffney, M. B. G. Hornillos, A. Görden, P. T. Greenlees, K. Helariutta, et al., *Phys. Rev. C* **81**, 014310 (2010).
- [15] R. D. Page, A. N. Andreyev, D. R. Wiseman, P. A. Butler, T. Grahn, P. T. Greenlees, R.-D. Herzberg, M. Huyse, G. D. Jones, P. M. Jones, et al., *Phys. Rev. C* **84**, 034308 (2011).
- [16] F. Iachello and A. Arima, *The interacting boson model* (Cambridge University Press, Cambridge, 1987).
- [17] A. Arima, T. Otsuka, F. Iachello, and I. Talmi, *Phys. Lett. B* **66**, 205 (1977).
- [18] T. Otsuka, A. Arima, F. Iachello, and I. Talmi, *Phys. Lett. B* **76**, 139 (1978).
- [19] T. Otsuka, A. Arima, and F. Iachello, *Nucl. Phys. A* **309**, 1 (1978).
- [20] P. D. Duval and B. R. Barrett, *Phys. Lett. B* **100**, 223 (1981).
- [21] P. D. Duval and B. R. Barrett, *Nucl. Phys. A* **376**, 213 (1982).
- [22] A. F. Barfield, B. R. Barrett, K. A. Sage, and P. D. Duval, *Z. Phys. A* **311**, 205 (1983).
- [23] C. De Coster, B. Decroix, K. Heyde, J. Jolie, H. Lehmann, and J. L. Wood, *Nucl. Phys. A* **651**, 31 (1999).
- [24] R. Fossion, K. Heyde, G. Thiamova, and P. Van Isacker, *Phys. Rev. C* **67**, 024306 (2003).
- [25] J. E. García-Ramos, V. Hellemans, and K. Heyde, *Phys. Rev. C* **84**, 014331 (2011).
- [26] A. Frank, P. Van Isacker, and C. E. Vargas, *Phys. Rev. C* **69**, 034323 (2004).
- [27] A. Frank, P. Van Isacker, and F. Iachello, *Phys. Rev. C* **73**, 061302 (2006).
- [28] I. O. Morales, A. Frank, C. E. Vargas, and P. Van Isacker, *Phys. Rev. C* **78**, 024303 (2008).
- [29] K. Heyde, C. De Coster, J. Jolie, and J. L. Wood, *Phys. Rev. C* **46**, 541 (1992).
- [30] C. De Coster, K. Heyde, B. Decroix, P. Van Isacker, J. Jolie, H. Lehmann, and J. L. Wood, *Nucl. Phys. A* **600**, 251 (1996).
- [31] H. Lehmann, J. Jolie, C. De Coster, B. Decroix, K. Heyde, and J. L. Wood, *Nucl. Phys. A* **621**, 767 (1997).
- [32] M. Bender, P.-H. Heenen, and P.-G. Reinhard, *Rev. Mod. Phys.* **75**, 121 (2003).
- [33] T. H. R. Skyrme, *Nucl. Phys.* **9**, 615 (1958).
- [34] D. Vautherin and D. M. Brink, *Phys. Rev. C* **5**, 626 (1972).
- [35] J. Decharge and M. Girod and D. Gogny, *Phys. Lett. B* **55**, 361 (1975).
- [36] D. Vretenar, A. Afanasjev, G. Lalazissis, and P. Ring, *Phys. Rep.* **409**, 101 (2005).
- [37] T. Nikšić, D. Vretenar, and P. Ring, *Prog. Part. Nucl. Phys.* **66**, 519 (2011).
- [38] W. Nazarewicz, *Phys. Lett. B* **305**, 195 (1993).
- [39] T. Duguet, M. Bender, P. Bonche, and P.-H. Heenen, *Phys. Lett. B* **559**, 201 (2003).
- [40] M. Bender, P. Bonche, T. Duguet, and P.-H. Heenen, *Phys. Rev. C* **69**, 064303 (2004).
- [41] J. P. Delaroche, M. Girod, G. Bastin, I. Deloncle, F. Hannachi, J. Libert, M. G. Porquet, C. Bourgeois, D. Hojman, P. Kilcher, et al., *Phys. Rev. C* **50**, 2332 (1994).
- [42] R. R. Chasman, J. L. Egido, and L. M. Robledo, *Phys. Lett. B* **513**, 325 (2001).
- [43] R. R. Rodríguez-Guzmán, J. L. Egido, and L. M. Robledo, *Phys. Rev. C* **69**, 054319 (2004).
- [44] J. L. Egido, L. M. Robledo, and R. R. Rodríguez-Guzmán, *Phys. Rev. Lett.* **93**, 082502 (2004).
- [45] T. Nikšić, D. Vretenar, P. Ring, and G. A. Lalazissis, *Phys. Rev. C* **65**, 054320 (2002).
- [46] K. Nomura, N. Shimizu, and T. Otsuka, *Phys. Rev. Lett.* **101**, 142501 (2008).
- [47] K. Nomura, T. Otsuka, N. Shimizu, and L. Guo, *Phys. Rev. C* **83**, 041302 (2011).
- [48] K. Nomura, N. Shimizu, D. Vretenar, T. Nikšić, and T. Otsuka, *Phys. Rev. Lett.* **108**, 132501 (2012).
- [49] K. Nomura, R. Rodríguez-Guzmán, L. M. Robledo, and N. Shimizu, *Phys. Rev. C* **86**, 034322 (2012).
- [50] S. Goriely, S. Hilaire, M. Girod, and S. Péru, *Phys. Rev. Lett.* **102**, 242501 (2009).
- [51] R. Rodríguez-Guzmán, P. Sarriguren, L. M. Robledo, and S. Perez-Martin, *Phys. Lett. B* **691**, 202 (2010).
- [52] R. Rodríguez-Guzmán, P. Sarriguren, and L. M. Robledo, *Phys. Rev. C* **82**, 044318 (2010).
- [53] R. Rodríguez-Guzmán, P. Sarriguren, and L. M. Robledo, *Phys. Rev. C* **82**, 061302 (2010).
- [54] J. F. Berger, M. Girod, and D. Gogny, *Nucl. Phys. A* **428**, 23 (1984).
- [55] A. Bohr and B. M. Mottelsson, *Nuclear Structure*, vol. 2 (Benjamin, New York, USA, 1975).
- [56] J. N. Ginocchio and M. W. Kirson, *Nucl. Phys. A* **350**,

- 31 (1980).
- [57] D. J. Thouless and J. G. Valatin, Nucl. Phys. **31**, 211 (1962).
- [58] H. Schaaser and D. M. Brink, Nucl. Phys. A **452**, 1 (1986).
- [59] K. Nomura, T. Otsuka, R. Rodríguez-Guzmán, L. M. Robledo, and P. Sarriguren, Phys. Rev. C **84**, 054316 (2011).
- [60] Brookhaven National Nuclear Data Center, <http://www.nndc.bnl.gov> (ENSDF).
- [61] D. D. Warner and R. F. Casten, Phys. Rev. C **28**, 1798 (1983).
- [62] A. S. Davydov and G. F. Filippov, Nucl. Phys. **8**, 237 (1958).